

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal62lcon

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	JAN 02	STN pricing information for 2008 now available
NEWS	3	JAN 16	CAS patent coverage enhanced to include exemplified prophetic substances
NEWS	4	JAN 28	USPATFULL, USPAT2, and USPATOLD enhanced with new custom IPC display formats
NEWS	5	JAN 28	MARPAT searching enhanced
NEWS	6	JAN 28	USGENE now provides USPTO sequence data within 3 days of publication
NEWS	7	JAN 28	TOXCENTER enhanced with reloaded MEDLINE segment
NEWS	8	JAN 28	MEDLINE and LMEDLINE reloaded with enhancements
NEWS	9	FEB 08	STN Express, Version 8.3, now available
NEWS	10	FEB 20	PCI now available as a replacement to DPCI
NEWS	11	FEB 25	IFIREF reloaded with enhancements
NEWS	12	FEB 25	IMSPRODUCT reloaded with enhancements
NEWS	13	FEB 29	WPINDEX/WPIDS/WPIX enhanced with ECLA and current U.S. National Patent Classification
NEWS	14	MAR 31	IFICDB, IFIPAT, and IFIUDB enhanced with new custom IPC display formats
NEWS	15	MAR 31	CAS REGISTRY enhanced with additional experimental spectra
NEWS	16	MAR 31	CA/CAPLUS and CASREACT patent number format for U.S. applications updated
NEWS	17	MAR 31	LPCI now available as a replacement to LDPCI
NEWS	18	MAR 31	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	19	APR 04	STN AnaVist, Version 1, to be discontinued
NEWS	20	APR 15	WPIDS, WPINDEX, and WPIX enhanced with new predefined hit display formats
NEWS	21	APR 28	EMBASE Controlled Term thesaurus enhanced
NEWS	22	APR 28	IMSRESEARCH reloaded with enhancements
NEWS EXPRESS	FEBRUARY 08 CURRENT WINDOWS VERSION IS V8.3, AND CURRENT DISCOVER FILE IS DATED 20 FEBRUARY 2008		
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS LOGIN	Welcome Banner and News Items		
NEWS IPC8	For general information regarding STN implementation of IPC 8		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN. Columbus * * * * *

FILE 'HOME' ENTERED AT 09:51:09 ON 08 MAY 2008

=> FILE CAPLUS

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 09:51:35 ON 08 MAY 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 8 May 2008 VOL 148 ISS 19

FILE LAST UPDATED: 7 May 2008 (20080507/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=>


Uploading C:\Program Files\Stnexp\Queries\GM1.str

L1 STRUCTURE UPLOADED

=> D L1

L1 HAS NO ANSWERS

L1 STR



G1 Cb,Cy,Hy

Structure attributes must be viewed using STN Express query preparation.

=> S L1 FULL

REGISTRY INITIATED

Substance data SEARCH and crossover from CAS REGISTRY in progress...

Use DISPLAY HITSTR (or FHITSTR) to directly view retrieved structures.

FULL SEARCH INITIATED 09:52:20 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 578 TO ITERATE

100.0% PROCESSED 578 ITERATIONS
 SEARCH TIME: 00.00.01

9 ANSWERS

L2 9 SEA SSS FUL L1

L3 12 L2

=> FILE CAPLUS

COST IN U.S. DOLLARS

SINCE FILE

ENTRY

TOTAL

SESSION

FULL ESTIMATED COST

0.48

179.53

FILE 'CAPLUS' ENTERED AT 09:52:52 ON 08 MAY 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 8 May 2008 VOL 148 ISS 19
FILE LAST UPDATED: 7 May 2008 (20080507/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> S L3
L4 12 L2

=> FILE REG	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.48	180.01

FILE 'REGISTRY' ENTERED AT 09:53:40 ON 08 MAY 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 7 MAY 2008 HIGHEST RN 1019993-29-3
DICTIONARY FILE UPDATES: 7 MAY 2008 HIGHEST RN 1019993-29-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2008.

Please note that search-term pricing does apply when conducting SmartSELECT searches.


REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>
Uploading C:\Program Files\Stnexp\Queries\GM1.str

L5 STRUCTURE UPLOADED

=> D L5
L5 HAS NO ANSWERS
L5 STR



G1 Cb,Cy,Hy

Structure attributes must be viewed using STN Express query preparation.

=> S L5 FULL

FULL SEARCH INITIATED 09:54:12 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 578 TO ITERATE

100.0% PROCESSED 578 ITERATIONS

9 ANSWERS

SEARCH TIME: 00.00.01

L6

9 SEA SSS FUL L5

=> FILE CAPLUS

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

178.36

358.37

FILE 'CAPLUS' ENTERED AT 09:54:23 ON 08 MAY 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 8 May 2008 VOL 148 ISS 19

FILE LAST UPDATED: 7 May 2008 (20080507/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.
They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> S L6

L7 12 L6

=> D L7 IBIB ABS HITSTR 1-12

L7 ANSWER 1 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:1393130 CAPLUS

DOCUMENT NUMBER: 148:417386

TITLE: Trans-4-Iodo, 4'-boranyl-chalcone induces antitumor activity against malignant glioma cell lines in vitro and in vivo

AUTHOR(S): Sasayama, Takashi; Tanaka, Kazuhiro; Mizukawa, Katsu; Kawamura, Atsufumi; Kondoh, Takeshi; Hosoda, Kohkichi; Kohmura, Eiji

CORPORATE SOURCE: Department of Neurosurgery, Kobe University Graduate School of Medicine, 7-5-1, Kusunoki-cho, Chuo-ku, Kobe, 650-0017, Japan

SOURCE: Journal of Neuro-Oncology (2007), 85(2), 123-132
CODEN: JNODD2; ISSN: 0167-594X

PUBLISHER: Springer

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Chalcones are considered the precursors of flavonoids and have been identified as interesting compds. with antitumor properties. Boronic-chalcone derivs. are more toxic to breast cancer cells compared to normal breast cells. Here, we studied the antitumor activities of trans-4-Iodo, 4'-boranyl-chalcone (TLBC), which is a boronic-chalcone derivative, in several glioma cell lines. TLBC showed a dose-dependent inhibition with inhibitory concentration 50% value in the μM range (5.5-25.5 μM) in various glioma cell lines. Flow cytometric and western blot assay demonstrated that TLBC induced apoptosis independent of changes to the tumor suppressor p53. This cytotoxic effect was the caspase-dependent manner. Also, TLBC lowered levels of anti-apoptotic Bcl-2 and/or Bcl-XL protein in several of the cell lines. To examine the antitumor effect of TLBC in vivo, we used a malignant glioma xenograft model. This result showed that in the mice treated with TLBC at 20 mg/kg, mean tumor volume was reduced by 43.9% ($P < 0.01$) in comparison with the control group. Immunohistochem. and western blot anal. showed that Bcl-2 protein levels were decreased and Bax protein levels were slightly increased in the tumors injected with 20 mg/kg TLBC compared with the control tumors. Therefore, we conclude that TLBC may be a potential chemotherapeutic agent for human glioma.

IT 562823-84-1


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(trans-4-Iodo, 4'-boranyl-chalcone induced cell cycle arrest, apoptosis and decreased antiapoptotic Bcl-2 as well as Bcl-XL protein expression in human glioblastoma cell)

RN 562823-84-1 CAPLUS

CN Boronic acid, B-[4-[(2E)-3-(4-iodophenyl)-1-oxo-2-propen-1-yl]phenyl]-
(CA INDEX NAME)

Double bond geometry as shown.



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:90998 CAPLUS

DOCUMENT NUMBER: 146:158818

TITLE: Quaternary nitrogen heterocyclic compounds for detecting aqueous monosaccharides in physiological fluids

INVENTOR(S): Geddes, Chris D.; Badugu, Ramachandram; Lakowicz, Joseph R.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 72pp., Cont.-in-part of Appl. No. PCT/US2004/022717.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070020182	A1	20070125	US 2005-318663	20051227
WO 2005000109	A2	20050106	WO 2004-US22717	20040628
WO 2005000109	A3	20050310		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2003-483124P P 20030627
US 2003-483202P P 20030627
WO 2004-US22717 A2 20040628

AB Quaternary nitrogen heterocyclic boronic acid-containing compds. are described, which are sensitive to glucose and fructose, as well as a variety of other physiol. important analytes, such as aqueous chloride and iodide, and a method of using the compds. Also disclosed is a contact lens doped with the quaternary nitrogen heterocyclic boronic acid-containing compound, and a method of using the doped contact lens to measure the concentration of analyte in tears under physiol. conditions.


IT 406719-92-4

RL: ARG (Analytical reagent use); PRP (Properties); ANST (Analytical study); USES (Uses)


(quaternary nitrogen heterocyclic compds. for detecting aqueous monosaccharides in physiol. fluids)

RN 406719-92-4 CAPLUS

CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



L7 ANSWER 3 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2006:315088 CAPLUS
DOCUMENT NUMBER: 145:290
TITLE: Anticancer activities of novel chalcone and bis-chalcone derivatives
AUTHOR(S): Modzelewska, Aneta; Pettit, Catherine; Achanta, Geetha; Davidson, Nancy E.; Huang, Peng; Khan, Saeed R.
CORPORATE SOURCE: Division of Chemical Therapeutics, Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, Baltimore, MD, 21231, USA
SOURCE: Bioorganic & Medicinal Chemistry (2006), 14(10), 3491-3495
CODEN: BMECEP; ISSN: 0968-0896
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 145:290
GI



I


AB A series of novel chalcones and bis-chalcones containing boronic acid moieties has been synthesized and evaluated for antitumor activity against the human breast cancer MDA-MB-231 (estrogen receptor-neg.) and MCF7 (estrogen receptor-pos.) cell lines and against two normal breast epithelial cell lines, MCF-10A and MCF-12A. These mols. inhibited the growth of the human breast cancer cell lines at low micromolar to nanomolar concns., with five of them showing preferential inhibition of the human breast cancer cell lines. Furthermore, bis-chalcone I exhibited a more potent inhibition of colon cancer cells expressing wild-type p53 than of an isogenic cell line that was p53-null.

IT 888203-69-8P 888203-70-1P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(anticancer activities of chalcone and bis-chalcone derivs.)


RN 888203-69-8 CAPLUS

CN Boronic acid, B-[4-[1-oxo-3-(2-thienyl)-2-propen-1-yl]phenyl]- (CA INDEX NAME)



RN 888203-70-1 CAPLUS

CN Boronic acid, B-[4-(3-benzo[b]thien-2-yl-1-oxo-2-propen-1-yl)phenyl]- (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:1114291 CAPLUS

DOCUMENT NUMBER: 145:58663

TITLE: A glucose-sensing contact lens: a new approach to noninvasive continuous physiological glucose monitoring

AUTHOR(S): Badugu, Ramachandram; Lakowicz, Joseph R.; Geddes, Chris D.

CORPORATE SOURCE: Cent. fluorescence Spectroscopy, Dep. Biochem. & Mol. Biol., Univ. of Maryland School of Medicine, MD, 21201, USA

SOURCE: Proceedings of SPIE-The International Society for Optical Engineering (2004), 5317(Optical Fibers and Sensors for Medical Applications IV), 234-245
CODEN: PSISDG; ISSN: 0277-786X

PUBLISHER: SPIE-The International Society for Optical Engineering


DOCUMENT TYPE: Journal

LANGUAGE: English

AB The authors have developed a new technol. for the non-invasive continuous monitoring of tear glucose using a daily use, disposable contact lens, embedded with sugar-sensing boronic acid containing fluorophores. The authors' findings show that the authors' approach may be suitable for the continuous monitoring of tear glucose levels in the range 50-500 μ M, which track blood glucose levels that are typically \approx 5-10-fold higher. The authors initially tested the sensing concept with well-established, previously published, boronic acid probes and the results could conclude the used probes, with higher pKa values, are almost insensitive toward glucose within the contact lens, attributed to the low pH and polarity inside the lens. Subsequently, the authors have developed a range of probes based on the quinolinium backbone, having considerably lower pKa values, which enables them to be suitable to sense the physiol. glucose in the acidic pH contact lens. Herein the authors describe the

results based on the authors' findings towards the development of glucose sensing contact lens and therefore an approach to non-invasive continuous monitoring of tear glucose using a contact lens.


IT 406719-92-4, Chalc 1
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES
(Uses)
(noninvasive continuous physiol. glucose monitoring in contact lens)
RN 406719-92-4 CAPLUS
CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



REFERENCE COUNT: 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 5 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2005:952799 CAPLUS
DOCUMENT NUMBER: 143:398794
TITLE: Monitoring the Effects of Antagonists on Protein-Protein Interactions with NMR Spectroscopy
AUTHOR(S): D'Silva, Loyola; Ozdowy, Przemyslaw; Krajewski, Marcin; Rothweiler, Ulli; Singh, Mahavir; Holak, Tad A.
CORPORATE SOURCE: Max Planck Institute for Biochemistry, Martinsried, D-82152, Germany
SOURCE: Journal of the American Chemical Society (2005), 127(38), 13220-13226
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB We describe an NMR method that directly monitors the influence of ligands on protein-protein interactions. For a two-protein interaction complex, the size of one component should be small enough (less than .apprx.15 kDa) to provide a good quality 15N (13C) HSQC spectrum after 15N(13C) labeling. The size of the second unlabeled component should be large enough so that the mol. weight of the preformed complex is larger than .apprx. 40 kDa. When the smaller protein binds to a larger one, broadening of NMR resonances results in the disappearance of most of its cross-peaks in the HSQC spectrum. Addition of an antagonist that can dissociate the complex would restore the HSQC spectrum of the smaller component. The method directly shows whether an antagonist releases proteins in their wild-type folded states or whether it induces their denaturation, partial unfolding, or precipitation We illustrate the method by studying lead compds. that have recently been reported to block the MDM2-p53 interaction. Activation of p53 in tumor cells by inhibiting its interaction with MDM2 offers new strategy for cancer therapy.
IT 562823-90-9
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(monitoring effects of antagonists on protein-protein interactions with NMR spectroscopy)
RN 562823-90-9 CAPLUS
CN Boronic acid, [4-[(2E)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 6 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:14124 CAPLUS

DOCUMENT NUMBER: 142:89411

TITLE: Quaternary nitrogen heterocyclic compounds for detecting aqueous monosaccharides in physiological fluids

INVENTOR(S): Geddes, Chris D.; Badugu, Ramachandran; Lakowitz, Joseph R.

PATENT ASSIGNEE(S): University of Maryland Biotechnology Institute, USA

SOURCE: PCT Int. Appl., 120 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005000109	A2	20050106	WO 2004-US22717	20040628
WO 2005000109	A3	20050310		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1644330	A2	20060412	EP 2004-778295	20040628
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
US 20070020182	A1	20070125	US 2005-318663	20051227
PRIORITY APPLN. INFO.:			US 2003-483124P	P 20030627
			US 2003-483202P	P 20030627
			WO 2004-US22717	W 20040628

OTHER SOURCE(S): MARPAT 142:89411

AB Disclosed are quaternary nitrogen heterocyclic boronic acid-containing compds. which are sensitive to glucose and fructose, as well as a variety of other physiol. important analytes, such as aqueous chloride and iodide, and a method of using the compds. Also disclosed is a contact lens doped with the quaternary nitrogen heterocyclic boronic acid-containing compound, and a method of using the doped contact lens to measure the concentration of analyte in tears

under physiol. conditions.


IT 406719-92-4

RL: ARG (Analytical reagent use); PRP (Properties); ANST (Analytical study); USES (Uses)

(quaternary nitrogen heterocyclic compds. for detecting aqueous monosaccharides in physiol. fluids)

RN 406719-92-4 CAPLUS

CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



L7 ANSWER 7 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:621102 CAPLUS

DOCUMENT NUMBER: 142:234524

TITLE: Cyanide-sensitive fluorescent probes

AUTHOR(S): Badugu, Ramachandram; Lakowicz, Joseph R.; Geddes, Chris D.

CORPORATE SOURCE: Center for Fluorescence Spectroscopy, Department of Biochemistry and Molecular Biology, Medical Biotechnology Center, University of Maryland School of Medicine, Baltimore, MD, 21201, USA

SOURCE: Dyes and Pigments (2005), 64(1), 49-55

CODEN: DYPIDX; ISSN: 0143-7208

PUBLISHER: Elsevier Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We characterize the response of several boronic acid containing fluorophores, which are widely used for sugar determination, towards aqueous cyanide. In two recent

reports we have shown that boronic acid containing fluorophores can be used to sense aqueous cyanide through physiol. safeguard levels. In this report we show that our new sensing mechanism is not just specific to our recently reported probes, but is indeed generic to the boronic acid moiety itself. Subsequently a wide range of cyanide-sensitive probes can now be realized, offering several modalities for fluorescence based cyanide sensing such as: intensity, lifetime, ratiometric, polarization and modulation fluorescence sensing.


IT 406719-92-4, Chalc 1

RL: ARG (Analytical reagent use); PRP (Properties); ANST (Analytical study); USES (Uses)

(cyanide sensing by boronic acid-containing fluorescent probes)

RN 406719-92-4 CAPLUS

CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



REFERENCE COUNT:

30

THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 8 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:1011575 CAPLUS

DOCUMENT NUMBER: 140:195805

TITLE: Noninvasive continuous monitoring of physiological glucose using a monosaccharide-sensing contact lens

AUTHOR(S): Badugu, Ramachandram; Lakowicz, Joseph R.; Geddes, Chris D.

CORPORATE SOURCE: Center for Fluorescence Spectroscopy, Department of Biochemistry and Molecular Biology, Medical Biotechnology Center, University of Maryland School of Medicine, Baltimore, MD, 21201, USA

SOURCE: Analytical Chemistry (2004), 76(3), 610-618

CODEN: ANCHAM; ISSN: 0003-2700

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We have tested the feasibility of tear glucose sensing using a daily, disposable contact lens embedded with boronic acid-containing fluorophores as a potential alternative to current invasive glucose-monitoring techniques. Our findings show that our approach may, indeed, be suitable for the continuous monitoring of tear glucose levels in the range 50-500 μ M, which track blood glucose levels that are .apprx.5-10-fold higher. We compare the response of the boronic acid probes in the contact lens to solution-based measurements and can conclude that both the pH and polarity within the contact lens need to be considered with respect to choosing/designing and optimizing glucose-sensing probes for contact lenses.


IT 661459-48-9

RL: ARU (Analytical role, unclassified); ANST (Analytical study)
(noninvasive continuous monitoring of physiol. glucose using monosaccharide-sensing contact lens)

RN 661459-48-9 CAPLUS

CN Boronic acid, [4-[(2E)-3-[4-(dimethylamino)phenyl]-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 9 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:1006923 CAPLUS

DOCUMENT NUMBER: 140:59511

TITLE: Preparation of boronic chalcone derivatives as anticancer agents

INVENTOR(S): Khan, Saeed R.

PATENT ASSIGNEE(S): Johns Hopkins University, USA

SOURCE: PCT Int. Appl., 56 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003106384	A2	20031224	WO 2003-US18962	20030612
WO 2003106384	A3	20040617		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003243594	A1	20031231	AU 2003-243594	20030612
US 20050176988	A1	20050811	US 2005-517781	20050420
PRIORITY APPLN. INFO.:			US 2002-388255P	P 20020613
			US 2003-444429P	P 20030203
			WO 2003-US18962	W 20030612
OTHER SOURCE(S):		MARPAT 140:59511		
GI				



AB The present invention relates to novel boronic chalcone derivs. I [Ar = (un)substituted heteroaryl, etc.; W = H, etc.; X = Zn, etc.; n = 0 or any integer; Z = (un)substituted alkylene, etc.] which are useful as antitumor/anticancer agents. The activity of compds. of this invention against the growth of human breast cancer cell lines was demonstrated.

IT 562823-84-1P 562823-90-9P 562823-91-0P
562823-92-1P 562823-93-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of boronic chalcone derivs. as anticancer agents)

RN 562823-84-1 CAPLUS

CN Boronic acid, B-[4-[(2E)-3-(4-iodophenyl)-1-oxo-2-propen-1-yl]phenyl]-(CA INDEX NAME)


Double bond geometry as shown.



RN 562823-90-9 CAPLUS

CN Boronic acid, [4-[(2E)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]phenyl]-(9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 562823-91-0 CAPLUS

CN Boronic acid, [4-[(2E)-3-(3-chloro-4-fluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 562823-92-1 CAPLUS

CN Boronic acid, [4-[(2E)-3-(3,4-difluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 562823-93-2 CAPLUS

CN Boronic acid, [4-[(2E)-3-(3-bromo-4-fluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



L7 ANSWER 10 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 2003:410905 CAPLUS
 DOCUMENT NUMBER: 139:117464
 TITLE: Design, Synthesis, and Evaluation of Novel
 Boronic-Chalcone Derivatives as Antitumor Agents
 AUTHOR(S): Kumar, Srinivas K.; Hager, Erin; Pettit, Catherine;
 Gurulingappa, Hallur; Davidson, Nancy E.; Khan, Saeed
 R.
 CORPORATE SOURCE: Division of Experimental Therapeutics, Sidney Kimmel
 Comprehensive Cancer Center at Johns Hopkins,
 Baltimore, MD, 21231, USA
 SOURCE: Journal of Medicinal Chemistry (2003), 46(14),
 2813-2815
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 139:117464


AB A series of boronic-chalcone derivs., e.g. 4-IC₆H₄CH:CHCOC₆H₄B(OH)₂-4,
 were synthesized and tested for antitumor activity against human breast
 cancer cell lines. The results show the boronic-chalcones are more toxic
 to breast cancer cells compared to normal breast cells than other known
 chalcones.
 IT 562823-84-1P 562823-90-9P 562823-91-0P
 562823-92-1P 562823-93-2P
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation);
 BIOL (Biological study); PREP (Preparation)
 (design, synthesis, and evaluation of novel boronic-chalcone derivs. as
 antitumor agents)
 RN 562823-84-1 CAPLUS
 CN Boronic acid, B-[4-[(2E)-3-(4-iodophenyl)-1-oxo-2-propen-1-yl]phenyl]-
 (CA INDEX NAME)

Double bond geometry as shown.




RN 562823-90-9 CAPLUS
 CN Boronic acid, [4-[(2E)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]phenyl]-
 (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 562823-91-0 CAPLUS
CN Boronic acid, [4-[(2E)-3-(3-chloro-4-fluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 562823-92-1 CAPLUS
CN Boronic acid, [4-[(2E)-3-(3,4-difluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



RN 562823-93-2 CAPLUS
CN Boronic acid, [4-[(2E)-3-(3-bromo-4-fluorophenyl)-1-oxo-2-propenyl]phenyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 11 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2002:211081 CAPLUS
DOCUMENT NUMBER: 137:134122
TITLE: Chalcone-analogue fluorescent probes for saccharides signaling using the boronic acid group
AUTHOR(S): DiCesare, Nicolas; Lakowicz, Joseph R.
CORPORATE SOURCE: Center for Fluorescence Spectroscopy, School of Medicine, University of Maryland, Baltimore, MD, 21201, USA
SOURCE: Tetrahedron Letters (2002), 43(14), 2615-2618

CODEN: TELEAY; ISSN: 0040-4039

PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English


AB Two new fluorescent probes based on 1,3-diphenylprop-2-en-1-one and on 1,5-diphenylpenta-2,4-dien-1-one structures are presented. Both probes possess one electron-donating dimethylamino group and one boronic acid group (electron-withdrawing group). The change between the neutral and the anionic form of the boronic acid group induced at high pH and/or in presence of sugar, induces optical changes for both probes. Spectroscopic data, pKa and dissociation consts. for different monosaccharides are presented and discussed in terms of sugar detection.

IT 406719-92-4

RL: ARU (Analytical role, unclassified); DEV (Device component use); PRP (Properties); ANST (Analytical study); USES (Uses)
(chalcone-analog fluorescent probes for saccharides signaling using the boronic acid group)

RN 406719-92-4 CAPLUS

CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 12 OF 12 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:73583 CAPLUS

DOCUMENT NUMBER: 136:275539

TITLE: New sensitive and selective fluorescent probes for fluoride using boronic acids

AUTHOR(S): DiCesare, Nicolas; Lakowicz, Joseph R.

CORPORATE SOURCE: Center for Fluorescence Spectroscopy, University of Maryland, School of Medicine, Baltimore, MD, 21201, USA

SOURCE: Analytical Biochemistry (2002), 301(1), 111-116

CODEN: ANBCA2; ISSN: 0003-2697

PUBLISHER: Academic Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We report the spectroscopic characterization of six fluorescent probes for fluoride sensing and/or monitoring. All probes are based on the ability of the boronic acid group to interact with fluoride. The probes combine electron donor and withdrawing groups and involve the excited charge transfer mechanism. The change between the neutral form of the boronic acid group [R-B(OH)2], which is an electron withdrawing group, and the anionic trifluoro form [R-BF3], which is an electron donating group, is at the origin of the different spectral changes observed for the investigated probes. Two probes are based on the stilbene structure where the boronic group in the 4 position is coupled with a cyano group, in one case, and the dimethylamino group in the other case, both at the 4' position. Another probe is based on the diphenyl-1,4-butadiene possessing the boronic acid group in the 4' position and a dimethylamino group in the 4' position. One probe is based on the diphenyloxazole structure having both the boronic acid and the dimethylamino groups in para positions. The two last probes reported are based on the benzalacetophenone (chalcone) structure, again coupling the boronic acid and dimethylamino groups. All


probes show spectral shifts and/or intensity changes in the presence of fluoride resulting in most of the cases to a wavelength-ratiometric way for the detection and/or anal. of fluoride. Selectivity and stability consts. are also presented and discussed. (c) 2002 Academic Press.

IT 406719-92-4

RL: ARU (Analytical role, unclassified); ANST (Analytical study)
(fluorescent probes for fluoride using boronic acids)

RN 406719-92-4 CAPLUS

CN Boronic acid, B-[4-[3-[4-(dimethylamino)phenyl]-1-oxo-2-propen-1-yl]phenyl]- (CA INDEX NAME)



REFERENCE COUNT:

28

THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT